

High-gradient operators in the N -vector model

S.E. Derkachov¹

Department of Mathematics, St. Petersburg Technology Institute

S.K. Kehrein²

Institut für Theoretische Physik, Ruprecht-Karls-Universität Heidelberg,
Philosophenweg 19, D-69120 Heidelberg, Germany

A.N. Manashov¹

Department of Theoretical Physics, State University St. Petersburg

February 1, 2008

Abstract

It has been shown by several authors that a certain class of composite operators with many fields and gradients endangers the stability of nontrivial fixed points in $2 + \epsilon$ expansions for various models. This problem is so far unresolved. We investigate it in the N -vector model in an $1/N$ -expansion. By establishing an asymptotic naive addition law for anomalous dimensions we demonstrate that the first orders in the $2 + \epsilon$ expansion can lead to erroneous interpretations for high-gradient operators. While this makes us cautious against over-interpreting such expansions (either $2 + \epsilon$ or $1/N$), the stability problem in the N -vector model persists also in first order in $1/N$ below three dimensions.

¹E-mail: manashov@phim.niif.spb.su

²E-mail: kehrein@physik.uni-augsburg.de

New address: Institut für Physik der Universität Augsburg, Theoretische Physik III, Elektronische Korrelationen und Magnetismus, D-86135 Augsburg, Germany

1 Introduction

Problems with high-gradient operators were first observed by Kravtsov, Lerner and Yudson in the Q -matrix model in a $2 + \epsilon$ expansion [1, 2]. They noticed that a certain class of canonically irrelevant composite operators with $2s$ gradients acquires anomalous dimensions in first order in ϵ that grow with s^2 and that finally make an infinite number of these operators relevant for $s \rightarrow \infty$ and any fixed finite $\epsilon > 0$. Wegner et al. have shown that this behaviour occurs also in other $2 + \epsilon$ expansions in one-loop order: In the N -vector model [3], the unitary matrix model [4] and the orthogonal matrix model [5]. In fact this behaviour has been observed in all $2 + \epsilon$ expansions that have so far been investigated with respect to this question. Judging from these results one could argue that the conventional nontrivial fixed points of these models become unstable against an infinite number of high-gradient perturbations. This would have important consequences for the present understanding of many problems that rely on $2 + \epsilon$ expansions.

A natural starting point to investigate the role of high-gradient operators in more detail is the universality class of the N -vector model. Here various expansions are available: $2 + \epsilon$, $1/N$ and $4 - \epsilon$ expansions. In Ref. [3] Wegner found a composite operator with $2s$ gradients and the full scaling dimension

$$y = 2(1 - s) + \epsilon \left(1 + \frac{s(s-1)}{N-2} \right) + O(\epsilon^2) \quad (1.1)$$

in the $2 + \epsilon$ expansion of the nonlinear σ -model

$$S = \frac{1}{2T} \int d^d x (\partial_\mu \vec{\pi})^2. \quad (1.2)$$

Here $\vec{\pi} = (\pi_1, \dots, \pi_N)$ is the normalized field $\vec{\pi}^2 = 1$ of the nonlinear σ -model. Operators with $y > 0$, $y = 0$ and $y < 0$ are relevant, marginal and irrelevant, resp. Castilla and Chakravarty have extended the calculation to two-loop order [6, 7] and obtained the term in order ϵ^2

$$y = 2(1 - s) + \epsilon \left(1 + \frac{s(s-1)}{N-2} \right) + \frac{\epsilon^2 s^3}{(N-2)^2} \left(\frac{2}{3} + O(1/s) \right) + O(\epsilon^3). \quad (1.3)$$

This two-loop result makes the stability problem even worse. For $s \gg N/\epsilon$ Eq. (1.3) amounts to an infinite number of relevant operators. On the other hand it could be proven in Ref. [8] that no similar problem occurs in the $4 - \epsilon$ expansion of $(\vec{\phi}^2)^2$ -theory. In the $4 - \epsilon$ expansion all anomalous dimensions in order ϵ can only make composite operators more irrelevant, but never go in the “dangerous” relevant direction.

Therefore somehow the behaviour seems to bend over between $d = 2$ and $d = 4$ dimensions. The obvious tool to investigate this question is the $1/N$ -expansion for $2 < d < 4$. This is the programme of this paper. Along its lines two other interesting aspects of the N -vector model emerge. First of all it is shown how the complete spectrum of anomalous dimensions for a subalgebra of composite operators in order $1/N$ is encoded as a straightforward diagonalization problem. This formally solves part of a programme by Lang and Rühl [9]. Then an asymptotic naive addition law for anomalous dimensions is derived that has also been found in the $4 - \epsilon$ expansion [10, 11]. This leads to a “clustering” of anomalous dimensions. However, this property is lost in a $2 + \epsilon$ expansion. It will become apparent that this serves as a warning against using the first orders of the $2 + \epsilon$ expansion to make statements about the high-gradient limit $s \rightarrow \infty$.

Still unfortunately this is not the end of the story. From an older result by Vasil’ev and Stepanenko [12] we will see that the high-gradient problem persists for $2 < d < 3$ dimensions

also after adding up all order in ϵ in first order in $1/N$. From their work one can see that powers σ^s of the auxiliary field σ in the $1/N$ -expansion interpolate between the high-gradient operators in $2 + \epsilon$ and composite operators $(\vec{\phi}^2)^s$ in the $4 - \epsilon$ expansion. For some reason this simple fact has so far remained unnoticed and we also want to use this paper to clarify this point.

Our paper is organized as follows. In Sect. 2 we discuss the renormalization of a certain class of composite operators to first order in the $1/N$ -expansion. Eigenoperators and their critical exponents are encoded as a straightforward diagonalization problem similar to Ref. [8]. In Sect. 3 an asymptotic naive addition law of anomalous dimensions valid for a large number of gradients is found that has already been observed in the $4 - \epsilon$ expansion [10, 11]. However, it is shown that this structure remains unobserved in the $2 + \epsilon$ expansion which can lead to misleading interpretations. In Sect. 4 it is shown that results in first order in $1/N$ obtained by Vasil'ev and Stepanenko in Ref. [12] allow the interpolation of the high-gradient behaviour between two and four dimensions. Sect. 5 contains the conclusions and suggestions for future work to solve the stability problem.

2 The spectrum of anomalous dimensions in order $1/N$

2.1 The operator subalgebra $\mathcal{C}_{\text{sym}}$

In this paper we shall mainly deal with composite operators $\Psi_{i_1, i_2 \dots i_l}^{A_1, A_2 \dots A_n}(x)$ constructed from n fields ϕ_A ($A = 1, \dots, N$) and their derivatives which are symmetric and traceless both with respect to their spatial $O(d)$ and internal $O(N)$ indices. We denote this important class of composite operators by $\mathcal{C}_{\text{sym}}$, compare also Ref. [13].

It is convenient to introduce “scalar” composite operators

$$\Psi(x, u, t) \equiv \Psi_{i_1 \dots i_l}^{A_1 \dots A_n}(x) u^{i_1} \dots u^{i_l} t_{A_1} \dots t_{A_n}, \quad (2.1)$$

where u and t are d -dimensional and N -dimensional vectors, resp. The original composite operators $\Psi_{i_1, i_2 \dots i_l}^{A_1, A_2 \dots A_n}(x)$ can be reconstructed from $\Psi(x, u, t)$ by differentiation with respect to u and t .

Now $\Psi(x, u, t)$ can generally be expressed as

$$\Psi(x, u, t) = \psi\left(\frac{\partial}{\partial a_1}, \dots, \frac{\partial}{\partial a_n}\right) \Phi(x; u; t; a_1, \dots, a_n) \Big|_{a_i=0}, \quad (2.2)$$

where

$$\Phi(x; u; t; a_1, \dots, a_n) \equiv P(u, \partial_v) T(t, \partial_s) \prod_{m=1}^n s^{A_m} \phi_{A_m}(x + a_m v) \Big|_{v, s=0}. \quad (2.3)$$

Here P and T are projectors on the subspace of traceless and symmetric tensors. The action of P (resp. T) on the operator $\Psi(x, u, t)$ amounts to the elimination of all terms proportional to u^2 (resp. t^2). The explicit form of the projector P, T can be found in Ref. [14], but will not be used in the following.

In this manner Eq. (2.2) establishes a one to one correspondence between a composite operator $\Psi(x, u, t) \in \mathcal{C}_{\text{sym}}$ and a symmetric polynomial $\psi(z_1, \dots, z_n)$, which we call the coefficient function of the operator $\Psi(x, u, t)$.

Now we briefly discuss the important subclass of conformal operators in $\mathcal{C}_{\text{sym}}$ that will play a central role in our further investigations. The operator $\Psi_{i_1, i_2 \dots i_l}(x)$ (for a moment we omit

the internal indices) is called conformal if it transforms as a tensor under conformal coordinate transformations, i.e.

$$\delta_\alpha \Psi_{i_1, \dots, i_l}(x) = (\alpha \partial + \frac{\Delta_\Psi}{d} \partial \alpha) \Psi_{i_1, \dots, i_l}(x) + \sum_{k=1}^l \partial_{i_k} \alpha_s \Psi_{i_1, \dots, s, \dots, i_l}(x) \quad (2.4)$$

at

$$x \rightarrow x'(x) = x + \alpha(x), \quad \delta_\alpha \phi(x) \equiv (\alpha(x) \partial + \frac{\Delta}{d} \partial \alpha(x)) \phi(x),$$

with $\alpha_i(x) = a_i + \omega_{ik} x_k + \lambda x_i + (x^2 \delta_{ik} - 2x_i x_k) c_k$.

Here Δ and Δ_Ψ are the conformal dimensions of the elementary field $\phi(x)$ and the operator $\Psi_{i_1, \dots, i_l}(x)$, resp. Note that for conformal operators from $\mathcal{C}_{\text{sym}}$ one simply has $\Delta_\psi = (\text{scale dimension}) - (\text{number of spatial indices})$.

A straightforward calculation shows that the coefficient function ψ of conformal operators must satisfy the following equation

$$\sum_{m=1}^n (z_m \partial_{z_m}^2 + \Delta \partial_{z_m}) \psi(z_1, \dots, z_n) = 0. \quad (2.5)$$

To find all solutions of this equation let us define the one to one map $\psi \rightarrow \hat{\psi}$:

$$\hat{\psi}(z_1, \dots, z_n) = \mathbf{T} \psi(z_1, \dots, z_n) = \frac{1}{\Gamma^n(\Delta)} \int_0^\infty \prod_{m=1}^n dt_m t_m^{\Delta-1} e^{-t_m} \psi(t_1 z_1, \dots, t_n z_n). \quad (2.6)$$

For the function $\hat{\psi}$ Eq. (2.5) reads

$$\sum_{m=1}^n \partial_{z_m} \hat{\psi}(z_1, \dots, z_n) = 0, \quad (2.7)$$

that is $\hat{\psi}$ must be translationally invariant. Thus the space of conformal operators with n fields and l derivatives is isomorphic to the vector space $\hat{C}(n, l)$ of symmetric, homogeneous of degree l and translationally invariant polynomials of n variables. The dimension of $\hat{C}(n, l)$ can be easily calculated from the generating function $D(n, x)$ [13]

$$D(n, x) \equiv \sum_{l=0}^{\infty} \dim C(n, l) x^l = \frac{1}{\prod_{p=2}^n (1 - x^p)}. \quad (2.8)$$

2.2 Calculation of Feynman diagrams

To describe the renormalization of operators from $\mathcal{C}_{\text{sym}}$ in the nonlinear σ -model we follow the scheme developed in the papers [12, 15]. The regularized action for the theory under consideration has the form

$$S = -\frac{1}{2} \partial_\mu \phi^A \partial^\mu \phi_A - \frac{1}{2} \sigma K_\omega \sigma + \frac{1}{2} M^\omega \sigma \phi^A \phi_A + \frac{1}{2} M^{2\omega} \sigma K \sigma. \quad (2.9)$$

Here ω and M are regularization parameter and regularization mass, resp. The kernel K is determined from the requirement of the cancellation of self-energy diagram contributions of the sigma field (see Fig. 1); $K(x) = -N/2G_\phi^2(x)$, where $G_\phi(x)$ is the propagator of the ϕ^A field ($\langle \phi^A(x) \phi^B(y) \rangle = \delta^{AB} G_\phi(x)$) [15].

The regularized kernel K_ω is defined as $K_\omega(x) = K(x)x^{-2\omega}$. The explicit expressions for the propagators G_ϕ and $G_\sigma = K_\omega^{-1}$ in momentum space are

$$G_\phi(p) = p^{-2}, \quad G_\sigma(p) = Bp^{-2(\mu-2+\omega)}, \quad (2.10)$$

where $\mu \equiv d/2$ and $B = -2 \cdot 4^{\mu+\omega} \pi^\mu \Gamma(2\mu - 2 + \omega) / (N\Gamma^2(\mu - 1)\Gamma(2 - \mu - \omega))$.

In the calculation of Feynman diagrams the divergencies appear as poles in ω which therefore corresponds to ϵ in the dimensional regularization scheme. It should be noted that the nonlinear σ -model is not multiplicatively renormalizable [15]. In general its critical exponents cannot be expressed via the renormalization constants Z_i but must be determined (employing conformal invariance of the theory [15]) directly from the renormalized Green functions by looking at the long distance logarithmic contributions. However, in Ref. [12] it was shown that in order to find the critical exponents in first order in $1/N$ it is sufficient to calculate the divergent pole term of the Feynman diagrams. A detailed discussion of this point can be found in [12].

Thus to find the critical exponents in first order in $1/N$ we must extract the divergent part in the corresponding diagrams.

For the operators from $\mathcal{C}_{\text{sym}}$ only the diagram in Fig. 2 contributes in first order in $1/N$. Diagrams containing cycles of ϕ lines are zero due to the $O(N)$ tracelessness of the operators in question. Diagrams describing transitions $\partial_i \partial_k \rightarrow \delta_{ik} \sigma$ vanish due to the tracelessness in the spatial indices.

As usual one has to consider the renormalization of the set of mixing operators. Renormalized operators are defined as

$$[\Psi_r(\phi)]_R = \Psi_r(\phi) + \frac{1}{\omega} \sum_{r'} Q^{rr'} \Psi_{r'}(\phi). \quad (2.11)$$

The critical exponents are obtained from the mixing matrix Q via

$$x_{\Psi}^{rr'}(g) = n\gamma_\phi \delta^{rr'} + 2Q^{rr'}, \quad (2.12)$$

where n is the number of fields ϕ in the operator $\Psi_r(\phi)$ and $\gamma_\phi = \eta/2$ is the anomalous dimension of the elementary field ϕ with

$$\eta = \frac{4}{N} \frac{\Gamma(2\mu - 2)}{\Gamma^2(\mu - 1)\Gamma(2 - \mu)} \frac{2 - \mu}{\Gamma(\mu + 1)} + O(1/N^2). \quad (2.13)$$

It is convenient to fulfill the renormalization procedure in terms of the generating function (2.3). The analytic expression corresponding to the diagram shown in Fig. 2 is

$$e^{\imath(u, p_i a_i + p_k a_k)} \int \frac{d^{2\mu} l}{(2\pi)^{2\mu}} \frac{e^{\imath(a_i - a_k)(u, l)}}{l^{2(\mu-2+\omega)} (l + p_i)^2 (p_k - l)^2}, \quad (2.14)$$

where $(\cdot, \cdot)$ denotes the scalar product. The calculation of this integral does not cause any problems and we give the final answer for the renormalization of the generating function at once

$$[\Phi(x; u; t; a_1, \dots, a_n)]_R = \Phi(x; u; t; a_1, \dots, a_n) + \frac{\nu(\mu)}{\omega N} \sum_{i < k} H_{ik} \Phi(x; u; t; a_1, \dots, a_n), \quad (2.15)$$

where

$$\nu(\mu) = 2 \frac{\Gamma(2\mu - 2)}{\Gamma^2(\mu - 1)\Gamma(2 - \mu)\Gamma(\mu - 2)} \quad (2.16)$$

and

$$H_{ik}\Phi(..a_i, .., a_k, ..) = 1/2(h_{ik} + \bar{h}_{ik})\Phi(..a_i, .., a_k, ..) =$$

$$1/2\left(\int_0^1 \prod_{m=1}^3 d\alpha_m \delta\left(\sum_{n=1}^3 \alpha_n - 1\right) \alpha_2^{\mu-3} \Phi(.., \alpha_1 a_i + \bar{\alpha}_1 a_k, .., \alpha_3 a_k + \bar{\alpha}_3 a_i, ..) + (i \leftrightarrow k)\right) \quad (2.17)$$

with the notation $\bar{\alpha}_m \equiv (1 - \alpha_m)$. On the rhs of this integral equation the i -th and k -th arguments are replaced, the other arguments (denoted by dots) remain unchanged. The first term in the brackets in Eq. (2.17) determines the action of the operator h_{ik} on the function $\Phi(.., a_i, .., a_k, ..)$, while the representation for $\bar{h}_{ik}$ is obtained from the latter by interchanging $a_i \leftrightarrow a_k$. Although on symmetric functions h_{ik} and $\bar{h}_{ik}$ are identical, it will be useful to distinguish them in the sequel.

To avoid misleadings we stress that (2.15) should be understood as an equation for generating functions, i.e. as an equation for the coefficients in the power series expansion in a_i of the lhs and rhs expressions. Eq. (2.17) also requires some comments. Due to the factor $\alpha_2^{\mu-3}$ the integrand has a nonintegrable singularity and the integral in rhs of Eq. (2.17) must be understood as the analytical continuation from the region $\mu > 2$. As a matter of fact $\alpha_2^{\mu-3}$ is always multiplied by some polynomial in α_i and the integral can always be expressed in terms of Γ -functions.

2.3 Diagonalization problem

To obtain the spectrum of critical exponents we must diagonalize the mixing matrix Q , i.e. find the multiplicatively renormalized eigenoperators. Taking into account Eqs. (2.2), (2.11), (2.15) it is easy to derive the eigenvalue equation for Q . Indeed, using the Fourier representation for $\phi(x + a_i u)$ ($\phi(x) = \int d^2\mu q \phi(q) e^{iqx}$) one immediately finds that the operator $\Psi(x, u, t)$ is multiplicatively renormalized if its coefficient function satisfies the following equation

$$\lambda \psi(z_1, \dots, z_n) = \sum_{i < k} \tilde{H}_{ik} \psi(z_1, \dots, z_n) = 1/2 \sum_{i < k} (\tilde{h}_{ik} + \tilde{\bar{h}}_{ik}) \psi(z_1, \dots, z_n), \quad (2.18)$$

where $\tilde{h}_{ik}$ again has a two-particle form

$$\tilde{h}_{ik} \psi(..z_i ..z_k..) = \int_0^1 \prod_{m=1}^3 d\alpha_m \delta\left(\sum_{n=1}^3 \alpha_n - 1\right) \alpha_2^{\mu-3} \psi(.., \alpha_1 z_i + \bar{\alpha}_3 z_k, .., \bar{\alpha}_1 z_i + \alpha_3 z_k, ..) . \quad (2.19)$$

Again on the rhs of this equation the i -th and k -th arguments are replaced in the manner shown. The expression for $\tilde{\bar{h}}_{ik}$ is obtained from (2.19) by interchanging $z_i \leftrightarrow z_k$. The critical exponent of the corresponding eigenoperator is

$$x = n(\mu - 1) + l + n \frac{\eta}{2} + \frac{2\nu(\mu)}{N} \lambda + O(1/N^2) \quad (2.20)$$

corresponding to a full scaling dimension $y = d - x$. Here and in the sequel n is the number of fields ϕ and l is the number of derivatives in the eigenoperator. The function $\nu(\mu)$ has been defined in (2.16) and η was given in (2.13).

Eq. (2.18) has a number of nice properties. But the eigenvalue problem is more manageable if written in terms of the $\hat{\psi}(z_1 .. z_n)$ functions. Under the transformation (2.6) the lhs and rhs of (2.18) transform into

$$\lambda \hat{\psi}(z_1, \dots, z_n) = \sum_{i < k} H_{ik} \hat{\psi}(z_1, \dots, z_n) \equiv H \hat{\psi}(z_1, \dots, z_n) \quad (2.21)$$

with the operator H_{ik} defined in Eq. (2.17). Some technical details of this calculation are given in Appendix A.

Now it is easy to show that the symmetry group of H is $SL(2, C)$. We define the action of the $SL(2, C)$ group on functions $\hat{\psi}(z_1, \dots, z_n)$ in the following manner

$$S(g)\hat{\psi}(z_1, \dots, z_n) = \prod_{i=1}^n (cz_i + d)^{-\Delta} \hat{\psi}\left(\frac{az_1 + b}{cz_1 + d}, \dots, \frac{az_n + b}{cz_n + d}\right) \quad (2.22)$$

where $g \in SL(2, C)$, $g = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$ and $ad - bc = 1$. It is easy to check that H and $S(g)$ commute for all g with $\Delta = \mu - 1$ in our case. The group $SL(2)$ has three generators S , S_+ , S_- :

$$S = \sum_{i=1}^n (z_i \partial_{z_i} + \Delta/2), \quad S_- = - \sum_{i=1}^n \partial_{z_i}, \quad S_+ = \sum_{i=1}^n (z_i^2 \partial_{z_i} + \Delta z_i). \quad (2.23)$$

All these generators commute with H and obey the commutation relations

$$[S, S_-] = -S_-, \quad [S, S_+] = +S_+, \quad [S_+, S_-] = 2S. \quad (2.24)$$

Using Eqs. (2.24) it is straightforward to see that the eigenvalue problem for H can be restricted to the subspace of functions annihilated by S_- :

$$S_{ln}^- = \{\hat{\psi}(z_1, \dots, z_n) | S_- \hat{\psi} = 0; S \hat{\psi} = (1 + n\Delta/2) \hat{\psi}\}. \quad (2.25)$$

According to (2.7) the operators in S_- are just the conformal operators transforming like (2.4). Like in Ref. [8] we therefore observe that the spectrum of critical exponents is generated by the conformal operators. All other eigenfunctions of H are obtained by application of S_+ on the eigenfunctions in S_{ln}^- , ($l, n = 0, 1, \dots$). Thus we have to solve the eigenvalues problem (2.21) for the operator H on the space of functions $\hat{\psi}(z_1, \dots, z_n)$ that are symmetric, translation invariant and homogeneous polynomials of degree l in n variables.

Another very useful property of H is its hermiticity with respect to a suitable scalar product defined on the space of homogeneous polynomials as

$$\langle \hat{\psi}_1 | \hat{\psi}_2 \rangle \equiv \psi_1(\partial_{z_1}, \dots, \partial_{z_n}) \hat{\psi}_2(z_1, \dots, z_n) |_{z_i=0}. \quad (2.26)$$

Remember that ψ and $\hat{\psi}$ are related via transformation (2.6). To prove the hermicity of H with respect to the scalar product (2.26)

$$\langle \hat{\psi}_1 | H \hat{\psi}_2 \rangle = \langle H \hat{\psi}_1 | \hat{\psi}_2 \rangle, \quad (2.27)$$

it is sufficient to note that $\tilde{H}_{ik} \psi(z_1, \dots, z_n) \rightarrow H_{ik} \hat{\psi}(z_1, \dots, z_n)$ under the transformation (2.6).

An important immediate consequence of this hermicity is the reality of the spectrum of critical exponents, i.e. all anomalous dimensions of eigenoperators in $\mathcal{C}_{\text{sym}}$ are real numbers at least in first order in $1/N$.

2.4 Double expansions

A closed analytical solution of Eq. (2.21) seems not possible except for the trivial case $n = 2$ where the form of the eigenfunctions is already fixed by conditions imposed on the functions $\hat{\psi}(z_1, z_2)$ alone. Indeed there is only one symmetric translation invariant polynomial of degree $l = 2L$: $\hat{\psi}_l(z_1, z_2) = (z_1 - z_2)^l$. The corresponding eigenvalue of (2.21) is $\lambda_l = 1/((l + \mu - 2)(l + \mu - 1))$ leading to critical exponents (2.20) in agreement with known results [16].

There are two important cases when the situation is simplified and the problem becomes more tractable. Let us consider the eigenvalue problem in the limiting cases where $d = 2 + \epsilon$ or $d = 4 - \epsilon$, in other words we want to hold the leading terms in ϵ in the anomalous dimensions. This corresponds to a double expansion in $1/N$ and ϵ keeping the leading nontrivial terms in Eq. (2.20). The multiplier $\nu(\mu)$ is in both cases proportional to ϵ^2 , whereas λ determined from Eq. (2.21) has a simple pole in ϵ . Extracting the pole contribution from the expression $H\hat{\psi}(z_1, \dots, z_n)$ (defined by Eq. (2.17)) one obtains the following formulae for the critical exponents with $O(\epsilon^2)$ accuracy in $d = 2 + \epsilon$

$$x = n \frac{\epsilon}{2} \left(1 + \frac{1}{N}\right) + l + \frac{\epsilon}{2N} \lambda_{2+\epsilon} + O(\epsilon^2) + O(1/N^2) \quad (2.28)$$

and $d = 4 - \epsilon$ dimensions

$$x = n \left(1 - \frac{\epsilon}{2}\right) + l + \frac{\epsilon}{N} \lambda_{4-\epsilon} + O(\epsilon^2) + O(1/N^2) , \quad (2.29)$$

where $\lambda_{2+\epsilon}$ and $\lambda_{4-\epsilon}$ are given by the eigenvalue problems

$$\lambda_{2+\epsilon} \hat{\psi}(z_1, \dots, z_n) = \sum_{i < k} \left(\hat{\psi}(\dots, z_i, \dots, z_i, \dots) + \hat{\psi}(\dots, z_k, \dots, z_k, \dots) \right) \quad (2.30)$$

and

$$\lambda_{4-\epsilon} \hat{\psi}(z_1, \dots, z_n) = 2 \sum_{i < k} \int_0^1 d\alpha \hat{\psi}(\dots, z_i \alpha + (1 - \alpha) z_k, \dots, z_i \alpha + (1 - \alpha) z_k, \dots) . \quad (2.31)$$

Again the i -th and k -th arguments on the rhs of these equations are replaced in the specified manner.

These double expansions can be compared with known results, which serves as a consistency check of our calculations. Formula (2.31) has already been obtained in Ref. [13] in a $4 - \epsilon$ expansion. In fact, one can also check in a lengthy calculation that the term in order ϵ^2/N from Ref. [10] is in agreement with our $1/N$ -results.

In the $2 + \epsilon$ expansion the complete spectrum of critical exponents in order ϵ has been obtained by Wegner [3]. The solution of the eigenvalue problem (2.30) is straightforward and yields

$$\lambda_{2+\epsilon} = m(2n - m - 1), \quad m = 0 \dots n . \quad (2.32)$$

Here m is the maximum number of elementary fields without any gradients acting on them in any term of the respective eigenoperator. This is in agreement with the results of Wegner [3] restricted to our subalgebra $\mathcal{C}_{\text{sym}}$ as can be checked easily.

3 An asymptotic naive addition law for anomalous dimensions

3.1 Outline of the proof

Since we have the exact solution of the eigenvalue problem for operators from $\mathcal{C}_{\text{sym}}$ in first order in the $2 + \epsilon$ expansion, it is natural to try to understand how large the corrections from higher orders in ϵ are. Indeed, the calculation of critical exponents of high-gradients operators in $\mathcal{C}_{\text{sym}}$ will lead to the observation that an appropriate expansion parameter is not ϵ but some combination of ϵ and l . Later we will see that a suitable combination is e.g. $\epsilon \cdot \ln l$.

The $1/N$ -expansion, in which the higher orders in ϵ are summed up, is very suitable for this problem. However, due to very fast growth of the size of the mixing matrix with increasing l it

seems hardly possible to obtain the analytical solution of the eigenvalue problem Eq. (2.21). But the following interesting quantitative prediction for critical exponents of high-gradient operators can be done.

Theorem: Let Λ_n^l be the spectrum of $H = \sum_{i < k} H_{ik}$ restricted to the subspace $S_-^{n,l}$, i.e. the subspace of symmetric homogeneous polynomials of degree l in n variables. Then for any two eigenvalues $\lambda_1 \in \Lambda_{n_1}^{R_1}$ and $\lambda_2 \in \Lambda_{n_2}^{R_2}$ corresponding to different conformal eigenoperators the following statement holds: There is a number L such that for all¹ $l > L$ one can find an eigenvalue $\lambda_l \in \Lambda_{n_1+n_2}^l$ such that

$$|\lambda_l - \lambda_1 - \lambda_2| \leq C/l^\Delta . \quad (3.1)$$

Here C is a fixed constant and $\Delta = \mu - 1$.

Hence any two anomalous dimensions for composite operators with n_1 and n_2 elementary fields generate a limit point given by the sum of these anomalous dimensions in the spectrum of composite operators with $n = n_1 + n_2$ fields. The approach to this limit point is given by (3.1). Therefore Eq. (3.1) is an “asymptotic naive addition law” for anomalous dimensions. The limit point structure in the spectrum leads to a clustering of anomalous dimensions. For more details we refer the reader to Refs. [10, 17] where this property was discussed in the framework of the $4 - \epsilon$ expansion.

The general scheme of the proof of the theorem is the same as in the paper [17]. So here we will only discuss the main ideas and some technical modifications as compared to [17]. The full proof can then be constructed along the same lines as in Ref. [17].

To prove the inequality (3.1) it is sufficient to show that in the subspace $S_-^{l,n}$ there exists a vector $\hat{\Psi}_S(z_1, \dots, z_n)$ such that

$$|(H - \lambda_1 - \lambda_2)\hat{\Psi}_S|/|\hat{\Psi}_S| \leq C/l^\Delta . \quad (3.2)$$

Here, as usual, $|\hat{\Psi}_S|^2 = \langle \hat{\Psi}_S | \hat{\Psi}_S \rangle$.

We begin with the construction of the test vector $\hat{\Psi}_S$. If $\hat{\psi}_1(z_1, \dots, z_{n_1})$ and $\hat{\psi}_2(z_1, \dots, z_{n_2})$ are the eigenfunctions of H corresponding to the eigenvalues λ_1 and λ_2 , we define $\hat{\Psi}_S$ in the following way

$$\hat{\Psi}_S(z_1, \dots, z_n) = \mathbf{T} \Psi_S(z_1, \dots, z_n) = \mathbf{Sym}_{\{z_1, \dots, z_n\}} \mathbf{T} \Psi(z_1, \dots, z_n) , \quad (3.3)$$

where $\mathbf{Sym}$ denotes symmetrization with respect to $z_1, \dots, z_n$. The function $\Psi(z_1, \dots, z_n)$ reads

$$\Psi(x, y) = \sum_{k=0}^l c_k (x_1 + \dots + x_{n_1})^k (y_1 + \dots + y_{n_2})^{l-k} \psi_1(x) \psi_2(y) , \quad (3.4)$$

where $c_k = (-1)^{l-k} C_k^l \Gamma(S_1) \Gamma(S_2) / \Gamma(k + S_1) \Gamma(l - k + S_2)$, $S_i = n_i \Delta + 2R_i$ and C_k^l is the binomial coefficient. Also, for brevity, we use the shorthand notation $\psi_{1,2}(x) = \psi_{1,2}(x_1, \dots, x_{n_{1,2}})$ and $\Psi(x, y) = \Psi(x_1, \dots, x_{n_1}, y_1, \dots, y_{n_2})$.

A straightforward calculation shows that $\Psi(z)$ defined by Eq. (3.4) satisfies Eq. (2.5). Consequently, the functions $\hat{\Psi}(z)$, $\hat{\Psi}_S(z)$ are translation invariant. The degree of the polynomial $\hat{\Psi}_S(z)$ is obviously equal to $l + R_1 + R_2$.

¹ If λ_1 and λ_2 correspond to the same eigenoperator, this statement holds only for even $l > L$.

To prove the inequality (3.2) for $\widehat{\Psi}_S$ given by Eq. (3.3) one needs to calculate $||\widehat{\Psi}_S||$ and $|(H - \lambda_1 - \lambda_2)\widehat{\Psi}_S|$. To do this we write the functions Ψ_S and $\widehat{\Psi}_S$ in the following more suitable form

$$\Psi(x, y) = \psi_1(x)\psi_2(y)K(\partial_a, \partial_b)\exp\left\{a\left(\sum_{i=1}^{n_1}x_i\right)+b\left(\sum_{j=1}^{n_2}y_j\right)\right\}\Big|_{a,b=0}, \quad (3.5)$$

where $K(a, b) = \sum_{k=0}^l c_k a^k b^{l-k}$.

Respectively, applying the transformation (2.6) to Eq. (3.4) yields after some algebra

$$\widehat{\Psi}(x, y) = \int_0^1 d\alpha d\beta \int_0^1 Du Dv f(\alpha, n_1, R_1) f(\beta, n_2, R_2) [\bar{\alpha}(ux) - \bar{\beta}(vy)]^l \psi_1(ux) \psi_2(vy), \quad (3.6)$$

where $f(\alpha, n, R) = (1 - \alpha)^{S-R-1} \alpha^{R-1} \cdot \Gamma(S)/\Gamma(R) \Gamma^n(\Delta)$; $Du = \prod_{i=1}^{n_1} du_i u_i^{\Delta-1} \delta(\sum_i u_i - 1)$; $ux \equiv \sum_i u_i x_i$ and $\psi(ux) = \psi(u_1 x_1, \dots, u_{n_1} x_{n_1})$.

Now the calculation of the norm of ψ_S can be done in the following manner:

1. Using formulae (2.26), (3.3) and taking into account that the functions $\psi_{1,2}(z)$ are fully symmetric one obtains

$$||\psi_S^l||^2 = (C_{n_1+n_2}^{n_1})^{-1} \sum_{k=0}^{n_2} C_k^{n_1} C_k^{n_2} A_l^{(k)}, \quad (3.7)$$

where $A_l^{(k)} = \Psi(\partial_x, \partial_y) \widehat{\Psi}(y_1, \dots, y_k, x_{k+1}, \dots, x_{n_1}, x_1, \dots, x_k, y_{k+1}, \dots, y_{n_2})$. Without loss of generality we now assume $n_2 \leq n_1$.

2. Using formulae (3.5), (3.6) and the translation invariance of $\widehat{\Psi}$ one can obtain the following expression for $A_l^{(k)}$

$$A_l^{(k)} = N(l) \int_0^1 d\alpha d\beta \int_0^1 Du Dv f(\alpha, n_1, R_1) f(\beta, n_2, R_2) \mathcal{A}^l(\alpha, \beta, u, v) \mathcal{F}(\alpha, \beta, u, v), \quad (3.8)$$

where $\mathcal{A}(\alpha, \beta, u, v) = [\bar{\alpha}(\bar{U}_k - U_k) + \bar{\beta}(\bar{V}_k - V_k)]/2$; $U_k = \sum_{i=0}^k u_i$; $V_k = \sum_{i=0}^k v_i$; $N(l) = K(\partial_a, \partial_b)(a - b)^l|_{a=b=0}$ and

$$\mathcal{F}(\alpha, ..) = \sum_{i=0}^{R_1+R_2} C_i^l \mathcal{A}^{-i} \partial_a^i \psi_1(\partial_x - a\bar{\beta}v, \partial_x - a\bar{\alpha}u) \psi_2(\partial_y + a\bar{\alpha}u, \partial_y - a\bar{\beta}v) \psi_1(uy, ux) \psi_2(vx, vy) \Big|_{y=-1/2}^{x=1/2}.$$

Here $\psi_i(x, y)$ means $\psi_i(x_1, \dots, x_k, y_{k+1}, \dots, y_{n_i})$ and we mention again that $\bar{x} \equiv (1 - x)$.

3. All the dependence on l in expression (3.8) for $A_l^{(k)}$, except the trivial one contained in the multiplier $N(l)$ (we do not need its explicit form), is due to the factor $\mathcal{A}^l$ in the integral. It is evident that the main contributions in the integral come from the regions in which $|\mathcal{A}(\dots)| \simeq 1$:

$$\{\mathcal{A} \simeq 1 \iff \alpha \simeq \beta \simeq U_k \simeq V_k \simeq 0\}; \quad \{\mathcal{A} \simeq -1 \iff \alpha \simeq \beta \simeq \bar{U}_k \simeq \bar{V}_k \simeq 0\}.$$

Then after long but straightforward calculations one finds that the leading contributions in the integral (3.8) from these regions behave as $C_1^{(k)}/l^{2k\Delta}$ for $A \simeq 1$ and $C_{-1}^{(k)}/l^{(n_1+n_2-2k)\Delta}$ for $A \simeq -1$. The explicit expressions for the constants $C_{\pm 1}^{(k)}$ are

$$C_1^{(k)} = \frac{\Gamma(S_1)\Gamma(S_1)}{\Gamma^{n_1+n_2-2k}(\Delta)} \int [D_{k+1}^{n_1} u] \psi_1^2(0, u) \int [D_{k+1}^{n_2} v] \psi_2^2(0, v), \quad (3.9)$$

$$C_{-1}^{(k)} = (-1)^l \frac{\Gamma(S_1)\Gamma(S_1)}{\Gamma^{2k}(\Delta)} \int [D_1^k u] (\psi_1(u, 0) \psi_2(u, 0)) \int [D_1^k v] (\psi_1(v, 0) \psi_2(v, 0)), \quad (3.10)$$

where $[D_k^n u] \equiv \prod_{i=k}^n du_i u_i^{\Delta-1} \delta(\sum_k^n u_i - 1)$.

Thus one can see that the main contribution to the norm of $\hat{\Psi}_S$ is given by $A_l^{(k)}$ when $k = 0$ or $2k = n_1 + n_2$. It can be shown that in these cases the exact answer for $A_l^{(k)}$ can be obtained and it coincides with the answer for the leading term of the asymptotic behaviour of $A_l^{(k)}$. Put in another way, the functions $\hat{\Psi}(z)$ with arguments arranged in different ways are almost orthogonal to one another for large l .

Eventually the expression for the norm of $\hat{\Psi}_S$ is

$$\|\hat{\Psi}_S\|^2 = N(l)(C_{n_1}^{n_1+n_2})^{-1} \|\hat{\psi}_1\|^2 \|\hat{\psi}_2\|^2 (1 + (-1)^l \delta_{\psi_1, \psi_2}) (1 + O(1/l^\Delta)) . \quad (3.11)$$

Our next task is to obtain an estimate from above for $\mathcal{E}(l) \equiv \|(H - \lambda_1 - \lambda_2)\hat{\Psi}_S\|$. Instead of trying to derive the exact asymptotic behaviour for this quantity, we prefer to give a rough estimate. This, however, reproduces the true dependence on l but in a much shorter calculation.

1. First of all, taking into account that $H = \sum_{i < k} H_{ik}$ is symmetric under transpositions of its arguments and using Schwartz's inequality one gets

$$\mathcal{E}(l) = \|(H - \lambda_1 - \lambda_2)\hat{\Psi}_S\| \leq \|(H - \lambda_1 - \lambda_2)\hat{\Psi}\| . \quad (3.12)$$

Remember that $\hat{\Psi}(z) = \hat{\Psi}(z)(z_1 \cdots z_n)$.

2. It can be easily shown that $(z_1 + \dots z_n)\psi(z) \rightarrow S_+ \hat{\psi}(z)$ under transformation (2.6). Then taking advantage of the explicit form of $\Psi(z)$ (Eq. (3.4)) and the $SL(2, C)$ invariance of H one obtains: $\sum_{i < k}^{n_1} H_{ik} \hat{\Psi}(z) = \lambda_1 \hat{\Psi}(z)$ and $\sum_{n_2 < i < k} H_{ik} \hat{\Psi}(z) = \lambda_2 \hat{\Psi}(z)$. Taking into account the symmetry properties of $\hat{\Psi}(z)$ and again using Schwartz's inequality we find

$$\mathcal{E}(l) \leq n_1 n_2 < \hat{\Psi}(x, y) H^2(x_1, y_1) \hat{\Psi}(x, y) > . \quad (3.13)$$

It is possible to simplify Eq. (3.13) even more. Let us remember that the eigenvalues of $H_{ik} = 1/2(h_{ik} + \bar{h}_{ik})$ are $\lambda_l = [(l + \mu - 2)(l + \mu - 1)]^{-1}$ $l = 0, 2, 4, \dots$. One sees that for $1 < \mu < 2$ the only negative eigenvalue is λ_0 . All the other eigenvalues lie in the interval $(0, 1)$. Thus if P_0 is the projector onto the subspace of eigenfunctions of $H(x, y)$ with $\lambda = \lambda_0$ one has $< \hat{\Psi} | H(x_1, y_1) | \hat{\Psi} > \leq < \hat{\Psi} | (H - \lambda_0 P_0) | \hat{\Psi} > + \lambda_0^2 < \hat{\Psi} | P_0 | \hat{\Psi} > \leq < \hat{\Psi} | \bar{h}(x_1, y_1) | \hat{\Psi} > + \lambda_0(\lambda_0 - 1) < \hat{\Psi} | P_0 | \hat{\Psi} >$.

The projector P_0 can be written as

$$P_0 \hat{\psi}(x, y) = B^{-1}(\Delta, \Delta) \int_0^1 ds (s(1-s))^{\Delta-1} \hat{\psi}(sx + (1-s)y, sx + (1-s)y) . \quad (3.14)$$

Here $B(x, y)$ is Euler's beta function.

3. The calculation of $< \hat{\Psi} | P_0 | \hat{\Psi} >$ and $< \hat{\Psi} | \bar{h} | \hat{\Psi} >$ is carried out in a similar manner, so we consider the latter only. First of all it is useful to rewrite the expression (2.17) for $\bar{h}_{ik}$ in a well-defined form for $2 < d < 4$ (compare Eq. (2.17) and the comments to it).

$$\bar{h}_{ik} \hat{\psi}(x, y) = (\Delta - 1)^{-1} \int_0^1 ds dt t(1-t)^{\Delta-1} (\mu + t\partial_t) \hat{\psi}((1-st)x + sty, t\bar{s}x + (1-\bar{s}t)y) . \quad (3.15)$$

Furthermore, in full analogy with the derivation of Eq. (3.8) we obtain

$$\begin{aligned} < \hat{\Psi} | \bar{h}(x_1, y_1) | \hat{\Psi} > = \frac{N(l)}{\Delta - 1} \int_0^1 d\alpha d\beta ds dt Du Dv f(\alpha, n_1, R_1) f(\beta, n_2, R_2) \\ & \times t(1-t)^{\Delta-1} (\mu + t\partial_t) \mathcal{A}^l(\alpha, \beta, u, v, t, s) \mathcal{G}(\alpha, \beta, u, v, t, s) , \end{aligned} \quad (3.16)$$

where $\mathcal{A} = [\bar{\alpha}(1/2 - u_1 ts) + \bar{\beta}(1/2 - v_1 t\bar{s})]$. The function $\mathcal{G}$ is a polynomial of degree $R_1 + R_2$ and has a structure like $\mathcal{F}$ in Eq. (3.8) Again one concludes that the dominant contribution

to the integral comes from the region where $|\mathcal{A}| \simeq 1$. At first sight the terms arising from $t\partial_t$ in (3.16) give the leading term of the asymptotic behaviour as $l \rightarrow \infty$, since differentiation of $\mathcal{A}^l$ produces additional power of l . But these additional powers of l come in pair with “small variables” $\sim (1 - \mathcal{A})$ that lead to the observation that the “ μ -term” and the “ $t\partial_t$ -term” in Eq. (3.16) have the same order in l . Both these terms can be treated on an equal footing. We consider the first of them. The condition $1 - \mathcal{A} \leq \varepsilon$ leads to $\alpha \simeq \beta \simeq u_1 t s \simeq v_1 t \bar{s} \leq \varepsilon$. Then in this region $\mathcal{A} \simeq (1 - \alpha/2)(1 - \beta/2)(1 - u_1 t s - v_1 t \bar{s}) + O(\varepsilon^2)$. Function $\mathcal{G}$ in its turn can be bounded by $Const \cdot l^{R_1+R_2}$. Using this the integration with respect to $u_2, \dots, u_{n_1}, v_2, \dots, v_{n_2}$ and α, β can be performed easily giving a factor $l^{-(R_1+R_2)}$. Thus the integral under consideration can be bounded from above by the following expression

$$C(\Delta)N(l) \int_0^1 du dv ds dt t(1-t)^{\Delta-1} \bar{u}^{(n_1-1)\Delta-1} \bar{v}^{(n_2-1)\Delta-1} (1 - ust - vt\bar{s})^l.$$

After tedious but straightforward calculations one shows that the leading term of the asymptotic expansion of this integral behaves as $l^{-2\Delta}$. So the final answer for $\mathcal{E}(l)$ is

$$\mathcal{E}(l) \leq CN(l)l^{-2\Delta}. \quad (3.17)$$

It is clear that the inequalities (3.2), (3.1) now follow immediately from Eqs. (3.11), (3.17).

3.2 Failure of the limit point structure in the $2 + \epsilon$ expansion

From the results in the previous subsection it is clear that the approach to the limit point close to two dimensions is like $l^{-\epsilon/2}$, where l is the total number of gradients. Hence the behaviour is non-uniform for $d \in [2, 4]$ and only uniform for $d \in [2 + \delta, 4]$ with some fixed $\delta > 0$. A suitable expansion parameter for high-gradient operators close to two dimensions is therefore necessarily some combination of ϵ and l , e.g. the most straightforward choice is $\epsilon \cdot \ln l$.

Let us make this more explicit by looking at the spectrum of conformal operators with $n = 3$ fields. The anomalous dimensions obtained by diagonalizing (2.21) are plotted in Fig. 3. Let us concentrate on the curves approaching the limit curve labeled $\phi^2 \otimes \phi$ given by the (trivial) sum of anomalous dimensions of ϕ^A and $h_{A_1 A_2} \phi^{A_1} \phi^{A_2}$. Here $h_{A_1 A_2}$ is a symmetric and traceless tensor. First of all one can readily check that the approach to the limit curve as a function of l is in fact like $l^{-d/2+1}$, with even l approaching from above and odd l from below. The approach is clearly non-uniform in the interval $d \in [2, 4]$. Close to two dimensions the derivatives of the curves even diverge for $l \rightarrow \infty$, corresponding to diverging terms in order ϵ^2/N in the $2 + \epsilon$ expansion.

That the limit point structure cannot be observed in the $2 + \epsilon$ expansion is also apparent from the diagonalization of (2.30). We find the following possible eigenvalues for different numbers of fields n according to (2.32):

$$\begin{aligned} n = 1 & : & \lambda_{2+\epsilon} &= 0 \\ n = 2 & : & \lambda_{2+\epsilon} &= 0, 2 \\ n = 3 & : & \lambda_{2+\epsilon} &= 0, 4, 6 \end{aligned}$$

This of course agrees with the values in Fig. 3 for $d = 2$. The naive sum of the eigenvalues belonging to ϕ^A and $h_{A_1 A_2} \phi^{A_1} \phi^{A_2}$ with $n = 1, 2$ is $\lambda_{2+\epsilon} = 2$, but this value is not contained in the above list of possible eigenvalues with $n = 3$ fields. Similar observations can be made quite generally in the spectrum of the $2 + \epsilon$ expansion, i.e. the limit point structure (3.1) remains invisible.

4 Vasil'ev and Stepanenko's results re-examined

4.1 Interpolation of the high-gradient behaviour for $2 < d < 4$

In the operator subalgebra $\mathcal{C}_{\text{sym}}$ discussed so far we have seen that one must be cautious if one uses ϵ as an expansion parameter in $d = 2 + \epsilon$ for high-gradient operators. This can lead to wrong conclusions about the limit of a large number of gradients $l \rightarrow \infty$. Now, however, the problematic class of high-gradient operators giving rise to Eqs. (1.1), (1.3) does not belong to $\mathcal{C}_{\text{sym}}$. The structure of the respective eigenoperators in the $2 + \epsilon$ expansion is [3]

$$\sum_{k=0}^l \gamma_k (\partial_+ \vec{\pi} \cdot \partial_- \vec{\pi})^{2(l-k)} (\partial_+ \vec{\pi} \cdot \partial_+ \vec{\pi})^k (\partial_- \vec{\pi} \cdot \partial_- \vec{\pi})^k \quad (4.1)$$

with some coefficients γ_k ($\gamma_0 \neq 0$) and $\partial_{\pm} = \frac{1}{\sqrt{2}}(\partial_x \mp i\partial_y)$. Hence these eigenoperators of the nonlinear σ -model (1.2) are $O(N)$ and $O(d=2)$ -scalar.

In this section we use the opportunity to show that the operators (4.1) have in fact already been investigated in the first order of the $1/N$ -expansion for $2 < d < 4$ by Vasil'ev and Stepanenko [12]. As this is not immediately apparent from their work, we want to clarify this point here.

In Ref. [12] Vasil'ev and Stepanenko calculated the critical exponents of the composite operator σ^s made up of powers of the auxiliary field σ from (2.9).² They find the following full scaling dimension

$$y = d - 2s - \eta s(d-1) \frac{(s-1)d(d-3) - 2(d-1)}{4-d} + O(N^{-2}) \quad (4.2)$$

with η from (2.13). If one expands this additionally in $d = 2 + \epsilon$ one finds

$$y = d - 2s + \frac{\epsilon}{N} s(s-1) + O(\epsilon^2) + O(N^{-2}) \quad (4.3)$$

or alternatively in $d = 4 - \epsilon$

$$y = d - 2s - 6 \frac{\epsilon}{N} s(s-2) + O(\epsilon^2) + O(N^{-2}). \quad (4.4)$$

One readily observes that (4.3) corresponds to the high-gradient operators (4.1) as the critical exponent is consistent with (1.1). In $d = 4 - \epsilon$ Eq. (4.4) corresponds to the composite operators $(\vec{\phi}^2)^s$ with the full scaling dimension [18]

$$y = d - 2s - \epsilon \frac{6s(s-2)}{N+8} + O(\epsilon^2). \quad (4.5)$$

As these critical exponents are unique in the respective expansions, this proves that powers of the auxiliary field σ^s interpolate between the high-gradient operators (4.1) with $2s$ fields and $2s$ gradients in $d = 2 + \epsilon$ and the composite operators $(\vec{\phi}^2)^s$ with $2s$ fields and no gradients in $d = 4 - \epsilon$. This might appear surprising at first sight as both σ^s and $(\vec{\phi}^2)^s$ contain no gradients. However, the canonical dimension of $\vec{\phi}$ is $x_{\phi} = d/2 - 1$, which varies as

$$\begin{aligned} d = 4 - \epsilon & : x_{\phi} = 1 - \epsilon/2 \\ d = 2 + \epsilon & : x_{\phi} = \epsilon/2 \end{aligned}$$

²In Ref. [12] the auxiliary field was denoted by ψ .

and hence allows to built gradients into the respective eigenoperators as d varies from four to two. Therefore the “most stable” eigenoperators $(\vec{\phi}^2)^s$ in $d = 4 - \epsilon$ are linked with the unstable high-gradient operators (4.1).

Eq. (4.2) allows to answer the question where this behaviour bends over between two and four dimensions. Eq. (4.2) can be written in the form

$$y = d - 2s - \frac{1}{N} \left(s(s-1) K(d) + O(s) \right) + O(N^{-2}) \quad (4.6)$$

with

$$K(d) = 4(d-3) \frac{\Gamma(d-2)}{\Gamma^2(d/2-1)\Gamma(2-d/2)\Gamma(d/2)} (d-1) . \quad (4.7)$$

$K(d)$ is plotted in Fig. 4. It has the following general behaviour:

$$\begin{aligned} K(d) &> 0 \quad \text{for } d > 3 \\ K(d) &= 0 \quad \text{for } d = 3 \\ K(d) &< 0 \quad \text{for } d < 3 \end{aligned}$$

As $K(d)$ multiplies the quadratic term in s in (4.6), this yields an infinite number of relevant operators in first order in $1/N$ for $2 < d < 3$ in the limit $s \gg N$. Hence $d = 3$ is the dimension where this seemingly unstable behaviour changes into the stable behaviour of the $4 - \epsilon$ expansions [8]. Of course this observation can be affected in any direction by the unknown terms in order $1/N^2$.

4.2 Contact exponents

As a short aside we briefly put the above observations into perspective of a scenario suggested by Duplantier and Ludwig [19]. They have argued that only field theories with negative contact exponents $\Theta \leq 0$ have the potential to describe multifractal scaling phenomena (leaving aside the use of analytical continuations like e.g. the replica method in field theory). A contact exponent [20] $\Theta = x_k - x_i - x_j$ is defined from the short-distance OPE

$$O_i(r_1) O_j(r_2) = |r_1 - r_2|^{x_k - x_i - x_j} O_k(r_2) + \text{subdominant terms with } x_{k'} > x_k. \quad (4.8)$$

Multifractal scaling requires [19] $\Theta \leq 0$, whereas in field theories one generally finds $\Theta \geq 0$ [19]. E.g. in the $4 - \epsilon$ expansion one has for the symmetric traceless part of $\phi^{A_1} \dots \phi^{A_n}$ the inequality $x_n - n x_1 = n(n-1)\epsilon/(N+8) + O(\epsilon^2) > 0$ in agreement with $\Theta \geq 0$. Similar for $(\vec{\phi}^2)^s$ one finds

$$x_{(\vec{\phi}^2)^s} - s x_{\vec{\phi}^2} = 6s(s-1) \frac{\epsilon}{N+8} + O(\epsilon^2) > 0 . \quad (4.9)$$

However, as we have seen in the previous subsection the behaviour (4.9) changes qualitatively below three dimensions in first order in $1/N$. For powers of the auxiliary field σ^s one deduces from Ref. [12]

$$\begin{aligned} x_{\sigma^s} - s x_{\sigma} &= \frac{1}{N} s(s-1) K(d) + O(N^{-2}) \\ &= \begin{cases} > 0 & \text{for } d > 3 \\ = 0 & \text{for } d = 3 \\ < 0 & \text{for } d < 3 \end{cases} \end{aligned} \quad (4.10)$$

with $K(d)$ defined in (4.7). Hence below three dimensions one finds the unusual behaviour for field theories

$$x_{\sigma^s} < s x_{\sigma} \text{ for } 2 < d < 3 \quad (4.11)$$

in first order in $1/N$ allowing for negative contact exponents $\Theta < 0$. In the framework of the $2 + \epsilon$ expansion this was already noticed in Ref. [19]. Notice that in Eq. (4.11) one is not forced to consider the limit $s \gg N$, (4.11) holds for arbitrary s and is in this sense a truly perturbative expansion in a possibly small parameter $\frac{s}{N}$.

Obviously, all this makes sense only if the usual nontrivial fixed point does not become unstable for $s \gg N$, $2 < d < 3$. Which cannot be answered conclusively for the time being as we have seen in the previous subsection.

5 Conclusions

In this paper we have investigated properties of high-gradient operators in the N -vector model. This class of operators shows a seemingly unstable behaviour due to the large corrections from anomalous dimensions (1.1), (1.3) found in Refs. [3, 6, 7]. Therefore it was the natural programme of our work to use an $1/N$ -expansion for $2 < d < 4$ dimensions to learn something about the behaviour of these operators as a function of d . Here we briefly sum up the main results.

For the subalgebra $\mathcal{C}_{\text{sym}}$ of composite operators introduced in Sect. 2.1 we have obtained the spectrum of critical exponents in first order in $1/N$ in a similar manner as in Refs. [8, 13, 10] for $4 - \epsilon$ expansions: The spectrum is encoded as a straightforward diagonalization problem (2.21) in the space of homogeneous, symmetric and translationally invariant polynomials. Let us mention that the spectrum of critical exponents in the $1/N$ -expansion has been the subject of various papers by Lang and Rühl [9] that aimed at finding algebraic structures in d dimensional conformal field theories. In $\mathcal{C}_{\text{sym}}$ our formalism gives the same results as theirs based on operators product expansions.

In Sect. 3 we proved a similar limit point structure (or: naive asymptotic addition law for anomalous dimensions) as seen in Refs. [10, 11] in $4 - \epsilon$ expansions: Two eigenoperators with anomalous dimensions λ_1, λ_2 and n_1, n_2 elementary fields generate the limit point $\lambda_1 + \lambda_2$ in the spectrum of conformal operators with $n_1 + n_2$ elementary fields. The approach to this limit point is like $l^{-d/2}$ where l is the total number of gradients. Such “limit curves” for anomalous dimensions (see Fig. 3) have already been observed in Ref. [16].³

In the $2 + \epsilon$ expansion, however, this limit point structure cannot be seen as the approach is like $l^{-\epsilon/2}$ if one uses ϵ as the expansion parameter (for a detailed discussion see Sect. 3.2). Unfortunately this observation could not solve the stability problem of $2 + \epsilon$ expansions as the problematic class of operators leading to (1.3) does not belong to $\mathcal{C}_{\text{sym}}$. Still, besides the interesting non-perturbative structure also its failure in the $2 + \epsilon$ expansion teaches the important lesson that ϵ might not be the suitable expansion parameter for high-gradient operators. E.g. one would also be tempted to read of a stability problem in order ϵ^2 in Fig. 3 close to $d = 2$ for curves with even l tending to the limit curve $\phi^2 \otimes \phi$. But this is bent over in higher order in ϵ as can be seen there since asymptotically the limit curve is reached. Generally one has to be careful with expansions like (1.1), (1.3) where higher orders are larger than previous orders for $s \gg \epsilon/N$.

The $O(N)$ and $O(d)$ -scalar high-gradient operators leading to (1.3) have in fact already been investigated in an $1/N$ -expansion by Vasil’ev and Stepanenko [12]. This remained unnoticed,

³As non-perturbative structures are always of interest in nontrivial field theories, our $1/N$ -results lead to the interesting conjecture that this limit structure holds to all orders in the N -vector model.

probably due to the fact that powers of the auxiliary field σ from (2.9) without any gradients turn out to interpolate between unstable high-gradient operators in $2 + \epsilon$ expansions and the unproblematic $(\phi^2)^s$ -operators in $4 - \epsilon$ expansions. This allowed the conclusion drawn from Eq. (4.6) that a stability problem for the nontrivial fixed point exists in first order in $1/N$ for $s \gg N$ below three dimensions.

Of course one can argue that higher orders in the $1/N$ -expansion will bend this behaviour over again, at least for some region of the d - N -plane. It is possible that the stability problem is only an artefact of the first order approximations and a proper treatment requires the investigation of the full perturbation series. This was just the situation for the limit point structure in the $2 + \epsilon$ expansion discussed above.

Considering the fact that other criticism of $2 + \epsilon$ or $1/N$ -expansions [21, 22] also remained unresolved and the importance of $2 + \epsilon$ expansions in general, the present situation must be considered unsatisfactory. One possible direction for future work could be along the lines of Dasgupta and Lau [23, 24]. Their conclusion that topological excitations are relevant in the low temperature phase of the Heisenberg model could be linked to the high-gradient problem [7].

Finally we pointed out in Sect. 4.2 that powers of the auxiliary field σ lead to negative contact exponents (4.11) for $2 < d < 3$. Negative contact exponents are very unusual in field theories and could be linked to multifractal scaling phenomena as argued in Ref. [19]. However, this interesting possibility relies on a stable nontrivial fixed point and hence a satisfactory solution of the above stability problem.

This work was supported by Grant 95-01-00569a of the Russian Fond for Fundamental Research and by INTAS Grant 93-2492-ext and is carried out under the research program of the International Center for Fundamental Physics in Moscow. S.K.K. acknowledges hospitality of the Theory Group at St. Petersburg State University where part of this work was performed.

Note added:

After completion of this work we were made aware of Ref. [25] where the role of topological defects in the $d = N = 3$ Heisenberg model has been re-examined using Monte Carlo simulations yielding different conclusions than Refs. [23, 24]. For recent work on topological defects in two dimensional systems see also Ref. [26].

Appendix A

Here we show that under transformation (2.6) Eq. (2.18) transforms into Eq. (2.21). It is obvious that it is sufficient to do this for the two particle case. For simplicity we consider the operator $\tilde{h}_{12}$ only

$$\tilde{h}_{12}\psi(z_1, z_2) = \psi_h(z_1, z_2) = \int_0^1 d\alpha \int_0^{1-\alpha} d\beta (1 - \alpha - \beta)^{\Delta-2} \psi(\beta z_1 + (1 - \alpha)z_2, \alpha z_2 + (1 - \beta)z_1) . \quad (\text{A.1})$$

Then from the definition of the $\hat{\psi}$ -function one has

$$\hat{\psi}_h(z_1, z_2) = \int_0^\infty dt_1 dt_2 t_1^{\Delta-1} t_2^{\Delta-1} \psi_h(z_1 t_1, z_2 t_2) e^{-t_1 - t_2} . \quad (\text{A.2})$$

After a change of variables ($t_1 + t_2 = T$, $t_1 = Ts$) we get

$$\begin{aligned} \hat{\psi}_h(z_1, z_2) = & \int_0^\infty dT T^{2\Delta-1} e^{-T} \int_0^1 d\gamma \int_0^1 d\alpha \int_0^{1-\alpha} d\beta (\gamma(1-\gamma))^{\Delta-1} \\ & \times (1-\alpha-\beta)^{\Delta-2} \psi(T(\beta\gamma z_1 + (1-\alpha)(1-\gamma)z_2), T(\alpha(1-\gamma)z_2 + (1-\beta)\gamma z_1)). \end{aligned} \quad (\text{A.3})$$

We perform the following change of variables in the integral:

$$\beta\gamma = \nu s; \quad (1-\gamma)(1-\alpha) = (1-s)\nu; \quad \alpha(1-\gamma) = t(1-\nu); \quad (1-\beta)\gamma = (1-t)(1-\nu) \quad (\text{A.4})$$

The last equation obviously follows from the three other equations. Then it is easy to derive the following relations:

$$0 \leq \nu \leq 1; \quad 0 \leq s + t \leq 1 \quad (\text{A.5})$$

and

$$\gamma(1-\gamma)d\alpha d\beta d\gamma = \nu(1-\nu)dv ds dt, \quad (\text{A.6})$$

$$(1-\alpha-\beta)^{\Delta-2}(\gamma(1-\gamma))^{\Delta-2} = (1-s-t)^{\Delta-2}(\nu(1-\nu))^{\Delta-2}. \quad (\text{A.7})$$

Combining everything one immediately finds

$$\hat{\psi}_h(z_1, z_2) = \int_0^1 ds \int_0^{1-s} dt (1-s-t)^{\Delta-2} \hat{\psi}(sz_1 + (1-s)z_2, tz_2 + (1-t)z_1) = h_{12} \hat{\psi}(z_1, z_2). \quad (\text{A.8})$$

References

- [1] V.E. Kravtsov, I.V. Lerner and V.I. Yudson, Zh. Eksp. Teor. Fiz. 94 (1988) 255; Sov. Phys. JETP 67 (1988) 1441
- [2] V.E. Kravtsov, I.V. Lerner and V.I. Yudson, Phys. Lett. A 134 (1989) 245
- [3] F. Wegner, Z. Phys. B 78 (1990) 33
- [4] F. Wegner, Nucl. Phys. B[FS] 354 (1991) 441
- [5] H. Mall and F. Wegner, Nucl. Phys. B[FS] 393 (1993) 495
- [6] G.E. Castilla and S. Chakravarty, Phys. Rev. Lett. 71 (1993) 384
- [7] G.E. Castilla and S. Chakravarty, *Is the phase transition in the Heisenberg model described by the $2 + \epsilon$ -expansion of the nonlinear σ -model?*, Preprint cond-mat/9605088 (1996)
- [8] S.K. Kehrein, F. Wegner and Y.M. Pismak, Nucl. Phys. B[FS] 402 (1993) 669
- [9] K. Lang and W. Rühl, Z. Phys. C 63 (1994) 531 and the references therein
- [10] S.K. Kehrein, Nucl. Phys. B[FS] 453 (1995) 777
- [11] S.E. Derkachov and A.N. Manashov, Nucl. Phys. B[FS] 455 (1995) 685
- [12] A.N. Vasil'ev and A.S. Stepanenko, Teor. Mat. Fiz. 95 (1993) 160; Theor. Math. Phys. 94 (1993) 471
- [13] S.K. Kehrein and F. Wegner, Nucl. Phys. B[FS] 424 (1994) 521
- [14] S.E. Derkachov and Yu.M. Pismak, J. Phys. A: Math. Gen. 26 (1993) 1419
- [15] A.N. Vasil'ev and M.Yu. Nalimov, Teor. Mat. Fiz. 55 (1983) 163; Theor. Math. Phys. 55 (1983) 423
- [16] K. Lang and W. Rühl, Nucl. Phys. B[FS] 400 (1993) 597
- [17] S.E. Derkachov and A.N. Manashov, *General scaling relation in the scalar ϕ^4 model*, Preprint SPbU-IP-96-8, hep-th/9604173, to appear in J. Phys. A: Math. Gen.
- [18] F. Wegner, Phys. Rev. B 6 (1972) 1891
- [19] B. Duplantier and A.W.W. Ludwig, Phys. Rev. Lett. 66 (1991) 247
- [20] J. des Cloizeaux, J. Phys. (Paris) 41 (1980) 223
- [21] J.L. Cardy and H.W. Hamber, Phys. Rev. Lett. 45 (1980) 499
- [22] H. Kunz and G. Zumbach, J. Phys. A: Math. Gen. 22 (1989) L1043
- [23] M-H. Lau and C. Dasgupta, J. Phys. A: Math. Gen. 21 (1988) L51
- [24] M-H. Lau and C. Dasgupta, Phys. Rev. B 39 (1989) 7212
- [25] C. Holm and W. Janke, J. Phys. A: Math. Gen. 27 (1994) 2553
- [26] G. Zumbach, Phys. Lett. A 200 (1995) 257

Figure captions

- Fig. 1. Cancellation of the self-energy diagram contribution of the σ -field.
- Fig. 2. Only diagram contributing in first order in $1/N$ for operator insertions from $\mathcal{C}_{\text{sym}}$.
- Fig. 3. The eigenvalues λ obtained by diagonalizing (2.21) in the subspace of conformal operators with $n = 3$ fields plotted as a function of $d = 2\mu$. The resp. critical exponents and anomalous dimensions can be obtained via Eq. (2.20). The number of gradients l is only given for the curves approaching the dashed limit curve $\phi^2 \otimes \phi$, the unlabelled curves approach other limit curves in agreement with our theorem. Notice that the normalization factor $2/\Gamma(d/2-2)$ has been chosen such that the values for $d = 2$ agree with the eigenvalues $\lambda_{2+\epsilon}$ in our definition (2.28), same for $d = 4$ and $\lambda_{4-\epsilon}$ using (2.29).
- Fig. 4. The function $K(d)$ from (4.6) plotted as a function of d . Observe the sign change in three dimensions separating stable from seemingly unstable behaviour.

$$\frac{1}{2} \text{---}\bigcirc\text{---} + \overset{\text{K}}{\text{---}\bullet\text{---}} = 0.$$





